

*

November 12, 2009

INFINITE TYPE GERMS OF SMOOTH HYPERSURFACES IN $\mathbb{C}^n$

JOHN ERIK FORNÆSS, LINA LEE, YUAN ZHANG

ABSTRACT. In this paper we discuss germs of smooth hypersurface in $\mathbb{C}^n$. We show that if a point on the boundary has infinite D'Angelo type, then there exists a formal complex curve in the hypersurface through that point.

1. INTRODUCTION

Let M be a $\mathcal{C}^\infty$ real hypersurface in $\mathbb{C}^n$ and $p \in M$. Then there is a $\mathcal{C}^\infty$ local defining function r of M near p , i.e., $M = \{r = 0\} \cap U$, where U is some neighborhood of p and $dr(p) \neq 0$. Let $\zeta : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^n, p)$ be a germ of a nonconstant holomorphic curve that sends 0 to p . Let $\nu(\zeta)$ denote the degree of vanishing at 0 of the function $|\zeta_1(t) - \zeta_1(0)| + \cdots + |\zeta_n(t) - \zeta_n(0)|$. Likewise, let $\nu(\zeta^*r)$ denote the order of vanishing of $r \circ \zeta$ at the origin.

Definition 1. The D'Angelo type of M at p , $\Delta(M, p)$ is defined as

$$\Delta(M, p) := \sup_{\zeta} \frac{\nu(\zeta^*r)}{\nu(\zeta)}.$$

We say that p is a point of finite type if $\Delta(M, p) < \infty$ and infinite type otherwise. We also denote

$$\Delta(M, p, \zeta) := \frac{\nu(\zeta^*r)}{\nu(\zeta)}.$$

In the convergent case, D'Angelo [1] proved the following theorem:

Theorem 1 (D'Angelo). *If M is a germ of a real analytic hypersurface in $\mathbb{C}^n$ and $p \in M$, then $\Delta(M, p) = \infty$ if and only if there exists a germ of a complex curve $\zeta(t) : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^n, p)$ lying on M and passing through p .*

In this paper we prove the formal analogue of this result. With a formal complex curve through p we mean an expression of the form $\zeta(t) = (\zeta_1(t), \dots, \zeta_n(t))$ where each ζ_j is a formal power series in the complex variable t and $\zeta(0) = p$.

Theorem 2. *If $M = \{r = 0\}$ is a germ of a $\mathcal{C}^\infty$ real hypersurface in $\mathbb{C}^n$ and $p \in M$, then $\Delta(M, p) = \infty$ if and only if there exists a nonconstant formal complex curve $\zeta(t) : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^n, p)$ such that $\zeta(0) = p$ and $r \circ \zeta$ vanishes to infinite order at 0.*

The proof is similar in outline to the proof by D'Angelo, but the formal analogue is somewhat different in some places. Denote by ${}_n\mathcal{O}_0$ the ring of germs of formal complex power series at 0 in $\mathbb{C}^n$. A key ingredient is the following

Theorem 3. *Let I be an ideal in ${}_n\mathcal{O}_0$. If $\dim({}_n\mathcal{O}_0/I) = \infty$, then there is a formal power curve ζ near 0 such that $\zeta^*f \sim 0$ for all $f \in I$.*

*The first author is supported by an NSF grant. Keywords: Plurisubharmonic Functions, D'Angelo type. 2000 AMS classification. Primary: 32T25; Secondary 32C25

In section 2 we collect background results. Then in Section 3 we prove the Theorem using Theorem 3, a version of the Nullstellensatz, and in the last section we prove Theorem 3. The authors are grateful to Steven Krantz for suggesting this problem.

2. PRELIMINARY RESULTS

2.1. Infinite matrices. Let $\mathcal{M}$ denote the set of infinite matrices

$$\begin{bmatrix} a_{11} & a_{12} & \dots \\ a_{21} & a_{22} & \dots \\ \dots & \dots & \dots \end{bmatrix}$$

where the $\{a_{ij}\}_{1 \leq i, j < \infty}$ are complex numbers.

For $A = \{a_{ij}\} \in \mathcal{M}$ let $A^* = \{\overline{a_{ji}}\}$ denote the adjoint matrix. We let $\mathcal{M}_0$ denote those matrices A for which all sums, $\sum_j |a_{ij}|^2 \leq 1, \sum_i |a_{ij}|^2 \leq 1$. If $A = \{a_{ij}\}, B = \{b_{k\ell}\} \in \mathcal{M}$ we define the matrix product $AB = \{c_{i\ell}\}$ where $c_{i\ell} = \sum_k a_{ik} b_{k\ell}$ if all these sums are convergent to a finite complex number. If $A, B \in \mathcal{M}_0$ then the matrix product $AB \in \mathcal{M}$ always exists. Also let $\mathcal{M}_1 \subset \mathcal{M}_0$ denote the unitary matrices, i. e. those for which $AA^* = A^*A = I =: \{\delta_{ij}\}$.

If $A = \{a_{ij}\} \in \mathcal{M}, A_k = \{a_{ij}^k\} \in \mathcal{M}_0, k = 1, \dots$ and for each $i, j, a_{ij}^k \rightarrow a_{ij}, k \rightarrow \infty$ we say that A_k converges weakly to A . In this case it is easy to see that $A \in \mathcal{M}_0$. In particular we note that if we have a sequence $U_k \in \mathcal{M}_0$, after taking a subsequence, we can assume that $U_k \rightarrow U \in \mathcal{M}_0$ and $U_k^* \rightarrow U^* \in \mathcal{M}_0$.

We let $\mathcal{M}_{1,k} \subset \mathcal{M}_1$ denote those unitary matrices $A = \{a_{ij}\}$ for which $a_{ij} = \delta_{ij}$ whenever $\max\{i, j\} > k$. Then the matrix ${}^kA = \{a_{ij}\}_{1 \leq i, j \leq k}$ is a unitary matrix on $\mathbb{C}^k$ and A looks like the identity matrix outside the first $k \times k$ block.

Let ℓ^2 denote the infinite sequences of complex numbers $c = \{c_j\}_{j \geq 1}$ with norm $\|c\|^2 = \sum_j |c_j|^2$. Then if $A \in \mathcal{M}_1$, then $\|Ac\| = \|c\|$. By weak convergence it follows that if $A \in \mathcal{M}_0$ is a weak limit of matrices in $\mathcal{M}_1$, then we still have $\|Ac\| \leq \|c\|$.

2.2. D'Angelo description of defining functions. Let r be the defining function of a smooth hypersurface around 0 in $\mathbb{C}^n$. For multiindices $J = (J_1, \dots, J_n), K = (K_1, \dots, K_n)$, denote $|J| := \sum_{j=1}^n J_j$ and set $J < K$ using some lexicographic ordering. In other words, we say $J < K$ if and only if either $|J| < |K|$, or $|J| = |K|$ and there exists $k \leq n$ such that $J_j = K_j$ for $j \leq k$ and $J_{k+1} < K_{k+1}$. Then we can write formally

$$\begin{aligned} r &\sim 2\Re h + 4\Re \sum_J \sum_{K \geq *J} a_{JK} z^J \bar{z}^K \\ &\sim 2\Re h + 4\Re \sum_J z^J \sum_{K \geq *J} a_{JK} \bar{z}^K \\ &\sim 2\Re h + \sum_J \left| z^J + \sum_{K \geq *J} \bar{a}_{JK} z^K \right|^2 - \sum_J \left| z^J - \sum_{K \geq *J} \bar{a}_{JK} z^K \right|^2 \\ &\sim \Re h + \sum_J |f_J|^2 - \sum_J |g_J|^2 \end{aligned}$$

where h is holomorphic, $f_J = z^J + \sum_{K \geq *J} \bar{a}_{JK} z^K$ and $g_J = z^J - \sum_{K \geq *J} \bar{a}_{JK} z^K$.

The key property for the above decomposition is that if we truncate r at any k only finitely many terms are nonzero.

3. FORMAL POWER SERIES CASE

For any given formal complex series g , we use $j_k(g)$ to denote the k -jets of g . A slight change of D'Angelo's theorem states as follows:

Lemma 1. *Let $\zeta = \zeta_k$ be a curve passing through the origin. If $j_{2k\nu(\zeta)}(\zeta^*r) = 0$, then there is an infinite unitary matrix U_k such that*

$$(1) \quad j_{2k\nu(\zeta)}(\zeta^*h) = j_{k\nu(\zeta)}(\zeta^*(f - U_k g)) = 0.$$

Proof. If $j_{2k\nu(\zeta)}(\zeta^*r) = 0$, then $j_{2k\nu(\zeta)}(\zeta^*h) = 0$ and $j_{2k\nu(\zeta)}(\|\zeta^*f\|^2) = j_{2k\nu(\zeta)}(\|\zeta^*g\|^2)$. Thus $\|j_{k\nu(\zeta)}\zeta^*f\|^2 = \|j_{k\nu(\zeta)}\zeta^*g\|^2$. Note the above norms make sense since there are only finite many terms in f and g with non-vanishing k -jets for each k according to our decomposition of r . For these finitely many f_J 's and g_J 's, applying ([1] Theorem 3.5), one can find an element $\tilde{U}$ of the group of unitary matrices such that

$$j_{k\nu(\zeta)}(\zeta^*(f - \tilde{U}g)) = 0.$$

Extending $\tilde{U}$ to an unitary operator U on l^2 by letting rest of the diagonal entries to be 1 and others 0, then we have (1) holds. $\square$

Denote by M_0 the maximal ideal in ${}_nO_0$ consisting of those which formally vanish at 0 and define $D(I) := \dim {}_nO_0/I$.

Lemma 2. *$D(I) < \infty$ if and only if for some integer k , $M_0^k \subset I$, where M_0 is a maximal ideal of ${}_nO_0$.*

Proof. If $M_0^k \subset I$, then $D(I) \leq D(M_0^k) < \infty$. If $D(I) = \ell < \infty$, then $z_j^\ell \in I$ for $j = 1, \dots, n$. Therefore $M_0^\ell \subset I$. $\square$

Lemma 3. *Suppose that the domain $r < 0$ has infinite type at 0. Then there exists a matrix $U \in \mathcal{M}_0$ which is a limit of matrices in $\mathcal{M}_1$ so that the ideal $I = I(h, f - Ug, U^*f - g)$ has infinite dimension.*

Proof. Since the domain has infinite type, there is a sequence of unitary matrices U_k as in the previous lemma. Let U be any weak limit of the U_k . Suppose that I has finite dimension. Then there exists an integer ℓ so that $M_0^\ell \subset I$. But then, for large enough k , $M_0^\ell \subset I(h, f - U_k g, U_k^*f - g)$, a contradiction. $\square$

Corollary 1. *Let $M = \{r = 0\}$ be a smooth hypersurface in $\mathbb{C}^n$. Let ζ be a formal curve. Then $\zeta^*r \sim 0$ if and only if there exists an operator $U \in \mathcal{M}_0$ which is a weak limit of operators in $\mathcal{M}_1$ such that $\zeta^*f \sim 0$ for any $f \in I$ as in Lemma 3.*

Proof. We only need to prove the sufficient direction, so consider any formal curve ζ such that $\zeta^*h \sim \zeta^*(f - Ug) \sim \zeta^*(U^*f - g) \sim 0$. Then $j_k(\zeta^*h) = 0$ and

$$\begin{aligned} \|j_k(\zeta^*f)\| &= \|j_k(\zeta^*Ug)\| = \|U(j_k(\zeta^*g))\| \\ &\leq \|j_k(\zeta^*g)\| \\ &= \|j_k(\zeta^*U^*f)\| = \|U^*(j_k(\zeta^*f))\| \\ &\leq \|j_k(\zeta^*f)\|. \end{aligned}$$

Therefore $\|j_k(\zeta^*f)\| = \|j_k(\zeta^*g)\|$ for any k . Letting k goes to infinity, then $\zeta^*h \sim \|\zeta^*f\|^2 - \|\zeta^*g\|^2 \sim 0$ and hence $\zeta^*r \sim 0$. $\square$

To complete the proof of Theorem 2, we only need to prove Theorem 3.

4. THEOREM 3

In the context of formal complex power series case, many parallel algebraic properties to those of convergent power series still hold. To our interest, we mention that the local ring ${}_nO_0$ is Noetherian. Moreover, Weierstrass preparation Theorem and division Theorem are both valid in the formal case. The reader may check [4], [5] etc for references.

For any proper ideal I in ${}_nO_0$, following the idea of Gunning([3]), we can similarly construct regular system of coordinates $z_1, \dots, z_n$ such that there exists an integer k satisfying:

1. ${}_kO_0 \cap I = 0$;
2. ${}_{j-1}O_0[z_j] \cap I$ contains a Weierstrass polynomial p_j in z_j for $j = k+1, \dots, n$.

In addition if I is prime, the theorem of the primitive element guarantees that by making a linear change of coordinates in $z_{k+1}, \dots, z_n$ plane, the quotient field ${}_n\tilde{M}_0$ of ${}_nO_0/I$ is an algebraic extension of ${}_kO_0$. i.e. ${}_n\tilde{M}_0 = {}_k\tilde{O}[\tilde{z}_{k+1}]$. We call the above regular system of coordinates to be *strictly regular system of coordinates*. Denote the discriminant of the unique irreducible defining polynomial p_{k+1} of z_{k+1} by D . Then for each coordinate function z_j , we can construct $q_j := D \cdot z_j - Q_j(z_{n+1}) \in I \cap {}_kO[z_{k+1}, z_j]$ for some $Q_j \in {}_kO[z_j]$, $j = k+2, \dots, n$. We also use $p_j \in {}_kO[z_j]$ to denote the defining Weierstrass polynomial of z_j . The ideal generated by $p_{k+1}, q_{k+2}, \dots, q_n$ is called to be the associated ideal.

The following lemma shows that the associated ideal $(p_{k+1}, q_{k+2}, \dots, q_n)$ and the original ideal cannot differ by very much.

Lemma 4. *There exists an integer ν such that*

$$D^\nu I \subseteq (p_{k+1}, q_{k+2}, \dots, q_n) \subseteq I.$$

If I is a principle ideal, we have the following result.

Lemma 5. *Let $P(z_1, \dots, z_k, w)$ be a formal Weierstrass polynomial $P = w^\ell + \sum_{j < \ell} b_j(z_1, \dots, z_k)w^j$, $b_j(0) = 0$. Suppose that $D(z_1, \dots, z_k) \neq 0$. Then there exists a formal curve ζ in $\mathbb{C}^{k+1}$ such that $P \circ \zeta \sim 0$.*

Proof. After linear change of coordinates on $(z_1, \dots, z_k)$, we assume $D(z_1, 0, \dots, 0) \neq 0$. Write $D(z_1, 0, \dots, 0) = \sum_{j=s}^{\infty} a_j z_1^j$ where $a_s \neq 0$. Next we consider truncations P_r of P where $r \gg s$. Then the discriminant of P_r , D_r as a polynomial in the elementary symmetric functions can still be written as $D_r(z_1, 0, \dots, 0) = a z_1^s + \dots$. For $z_1 \neq 0$ we can write $P_r(z_1, 0, \dots, 0) = \Pi_j (w - \alpha_j^r(z_1))$. The expression of D_r leads to the estimate $|\alpha_j^r - \alpha_i^r| \geq c|z_1|^s$ for some small c . Let $\Delta(\alpha_j^r(z_1), \epsilon|z_1|^s)$ be the disk with radius $\epsilon|z_1|^s$ centered at $\alpha_j^r(z_1)$ for each j . Then on the boundary of these discs $|P_r| \geq (\epsilon|z_1|^s)^\ell$. So for all higher truncations with $\rho > r \gg s\ell$ we have as well that $|P_\rho| \geq (\epsilon|z_1|^s)^\ell$ and that the zeroes of P_ρ are contained in these discs. We need to shrink z_1 for this to be true. We can also make these discs much smaller. This means that we have trapped the zero curves for P_r and so we can take formal limits and get a curve on which P vanishes to infinite order. $\square$

We have the following proposition:

Proposition 1. *Let $P \in {}_nO_0$ be a prime ideal with $D(P) = \infty$. Then there exists a formal curve ζ such that $\zeta^* f \sim 0$ for all $f \in P$.*

Proof. We first choose coordinates $z_1, \dots, z_n$ such that I is strictly regular. Note that since $D(P) = \infty$, $k \neq 0$.

Consider the associated ideals $(p_{k+1}, q_{k+2}, \dots, q_n)$. Notice the discriminant of p_{k+1} , D is not identically zero. Apply Lemma 5 to p_{k+1} , then there exists a formal power curve $\zeta' = (\zeta_1(t), \zeta_2(t), \dots, \zeta_{k+1}(t))$ such that $\zeta'^* p_{k+1} \sim 0$. We mention that the curve can always be chosen so that $D \circ (\zeta_1(t), \zeta_2(t), \dots, \zeta_k(t)) \neq 0$.

Next we need to add coordinates $z_{k+2}(t), \dots, z_n(t)$ so that all functions in the ideal vanish. Recall $q_j = Dz_j - Q_j(z_{k+1}) \in I$, $j > k+1$. Here $Q_j \in {}_nO[z_{k+1}]$. We solve for $z_j = Q_j/D$ and let $\zeta_j(t) = z_j$ for $j = k+2, \dots, n$. Then $(p_{k+1}, q_{k+2}, \dots, q_n)$ vanishes on $\zeta(t) = (\zeta_1(t), \zeta_2(t), \dots, \zeta_n(t))$. We'll show $z_j = Q_j/D$ is one of the formal roots for the defining Weierstrass polynomials $p_j \in {}_kO_0[z_j]$ and therefore Q_j/D vanishes at the origin. Indeed, for each $j > k+1$, denote by n_j the degree of $p_j \in {}_kO_0[z_j]$ with respect to z_j and consider $P_j = D^{n_j} p_j$. Replacing Dz_j in P_j by $q_j + Q_j$. We then get $P_j = H(q_j) + G(Q_j)$ for some polynomials $H(\cdot), G(\cdot)$ with no constant terms. Since $P_j \in I$ and p_{k+1} is the defining polynomial for z_{k+1} , $G(Q_j)$ is divisible by p_{k+1} . P_j therefore identically vanishes on the curve defined by $(z_1, \dots, z_{k+1}) = \zeta'(t)$, $q_j(\zeta') = 0$. $z_j = Q_j/D$ is then one of the formal roots for the defining Weierstrass polynomials p_j .

Finally we need to show for any f in the ideal I , f vanishes on the formal curve $\zeta(t)$. Indeed, by Lemma 4, there exists some high power ν , such that $D^\nu f \in (p_{k+1}, q_{k+2}, \dots, q_n)$. Therefore $(D^\nu f) \circ \zeta \sim 0$. Notice D vanishes to finite order on the curve, thus $\zeta^*(f) \sim 0$. $\square$

In order to prove Theorem 3, we also need the following simplification lemmas.

Lemma 6. *If $I = I_1 \cap I_2$ and $D(I) = \infty$, then $D(I_j) = \infty$ for some j .*

Proof. Suppose that both $D(I_j) < \infty$. Then for some integers k_1, k_2 we have that $M_0^{k_j} \subset I_j$; $j = 1, 2$. Hence $M_0^{\max\{k_1, k_2\}} \subset I$. Hence $D(I) < \infty$. $\square$

We also similarly define formal radical of an ideal I as follows

Definition 2. $\sqrt{I} := \{f : \text{there exists an integer } k \text{ such that } f^k \in I\}$.

Lemma 7. *If $D(P) = \infty$ then $D(\sqrt{P}) = \infty$.*

Proof. Suppose that $D(\sqrt{P}) < \infty$. Then for some k , $M_0^k \subset \sqrt{P}$. Let $f_1, \dots, f_s$ be generators of M_0^k . Then $f_i^{r_i} \in P$ for some r_i 's. If $f \in M_0^k$, then $f = \sum g_j f_j$, so $f^{r_1 + \dots + r_s} \in P$. Then $M_0^K \subset P$ for large enough K . But then $D(P) < \infty$. $\square$

Recall that an ideal J is primary if whenever $xy \in J$ then $x \in J$ and $y^n \in J$ for some n or vice versa.

We can now prove Theorem 3.

Let I be an ideal in ${}_nO_0$. If $D(I) = \infty$ then we want to show that there is a formal power curve ζ near 0 such that $\zeta^* f \sim 0$ for all $f \in I$.

Proof. By Lasker-Noether decomposition theorem, we can write $I = P_1 \cap \dots \cap P_k$, where P_j 's are primary ideals. By Lemma 6, we have $D(P_j) = \infty$ for some j . Lemma 7 also implies $D(\sqrt{P_j}) = \infty$. Notice $\sqrt{P_j}$ is prime, Proposition 1 implies the existence of a formal curve ζ such that $\zeta^* f \sim 0$ for all $f \in \sqrt{P_j}$. Since $I \subset P_j \subset \sqrt{P_j}$, $\zeta^* f \sim 0$ for all $f \in I$. The proof of theorem is thus complete. $\square$

REFERENCES

- [1] D'Angelo, J. P. Real hypersurfaces, orders of contact, and applications, *Annals of Math.* 115 (1982), 615-637.
- [2] D'Angelo, J. P. Several complex variables and the geometry of real hypersurfaces. *Studies in Advanced Mathematics*, 1992.
- [3] Gunning, R. C. Introduction to holomorphic functions of several variables. Volume II, local theory.
- [4] Lang, S. *Algebra*. 3rd ed. Addison Wesley. Revised third edition. *Graduate Texts in Mathematics*, 211. Springer-Verlag, New York, 2002.
- [5] Ruiz, J. The basic theory of power series. *Advanced Lectures in Mathematics*. Friedr. Vieweg and Sohn, Braunschweig, 1993.

John Erik Fornæss
Mathematics Department
The University of Michigan
East Hall, Ann Arbor, MI 48109
USA
fornaess@umich.edu

Lina Lee
Mathematics Department
The University of Michigan
East Hall, Ann Arbor, MI 48109
USA
linalee@umich.edu

Yuan Zhang
Mathematics Department
Rutgers University
New Brunswick, NJ, 08854
USA
yuanz@math.rutgers.edu